

Long-term neuroprotection of retinal ganglion cells by inhibiting caspase-2

Vigneswara, V; Ahmed, Z

DOI:

[10.1038/cddiscovery.2016.44](https://doi.org/10.1038/cddiscovery.2016.44)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Vigneswara, V & Ahmed, Z 2016, 'Long-term neuroprotection of retinal ganglion cells by inhibiting caspase-2', *Cell Death Discovery*, vol. 2, 16044. <https://doi.org/10.1038/cddiscovery.2016.44>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

CORRESPONDENCE

Long-term neuroprotection of retinal ganglion cells by inhibiting caspase-2

Cell Death Discovery (2016) 2, 16044; doi:10.1038/cddiscovery.2016.44;
published online 13 June 2016

Caspases are a family of cysteine-rich proteases that activate apoptotic pathways through their 'initiator' or 'executioner' pathways. Initiator caspases activate executioner caspase, such as caspase-2, -3, -6 and -7, which hydrolyze or cleave structural proteins in the cell leading to apoptosis. Caspase-2 is the most highly conserved caspase and uniquely fits into both initiator and executioner categories and as such partakes in a variety of apoptosis inducing signals including DNA damage, heat shock, endoplasmic reticulum and oxidative stress. Caspase-2-deficient neurons are resistant to apoptosis induced by beta-amyloid although it is involved in apoptosis of hippocampal neurons after ischemia and is expressed in ischemic retinae.¹

The protection of retinal ganglion cells (RGC), particularly after trauma and numerous ocular pathologies such as glaucoma and ischemic optic neuropathy, is of particular interest to the ocular biology field. RGC loss, a hallmark of these diseases leads to vision loss and blindness.² RGC are particularly susceptible and in the rat optic nerve transection/crush model in adult rats, RGC die rapidly and within 28 days, 90% of RGC are lost.³ As apoptosis accounts for the majority of RGC death, treatments that inhibit apoptosis afford limited neuroprotection, including microglial inhibitors, brain-derived neurotrophic factor, over-expression of Bcl-2 and caspase inhibitors.⁴ Caspase-3, -6, -8 and -9 are all regarded as important for RGC apoptosis, however, inhibition of these caspases, either alone or in specific combinations afforded protection of up to 60% of RGC.⁵ Furthermore, our observations that cleaved caspase-3 was absent in RGC and that little or no changes in cleaved caspase-6 and -7 were detected after transection of RGC axons by optic nerve crush (ONC) suggested that other caspases may have a part in RGC apoptosis.^{3,6}

We previously reported that RGCs exclusively expressed activated (cleaved) caspase-2 after ONC in adult rats and confirmed these findings by employing an unbiased caspase trapping assay.^{3,6} We further showed that inhibition of caspase-2 by either a pharmacological inhibitor⁶ or a chemically modified synthetic short interfering RNA (siCASP2) significantly protected RGC from death, the latter protecting >95% of RGC from death for up to 21 days after ONC.^{3,7} Here, we demonstrate in retinal whole mounts, prepared after FluroGold backfilling of RGC,^{3,7} that intravitreal injection of siCASP2 every 8 days protected RGC from death at all radial quadrants from the optic disc including the central, mid-periphery and outer quadrants, with >95% protection of RGC from death over the entire retina at 12 weeks after ONC (Supplementary Figure S1), compared with intact controls

(2170 ± 200 in siCASP2 treated eyes versus 2250 ± 230 RGC/mm² in intact controls). Only a small proportion of RGC were present in retinal whole mounts from control scrambled siRNA-(siCNL³) treated eyes (150 ± 130 RGC/mm²). Our results suggest that neuroprotection by siCASP2 is extremely efficient throughout the entire retina and that this is an unprecedented level of neuroprotection. To the best of our knowledge, our results far exceed those of any others published in the literature. Our results affirm that caspase-2 is the main caspase responsible for the death of RGC after direct trauma. This bodes well for siCASP2, otherwise known as QPI-1007, currently recruiting patients in a worldwide Phase III trial that is being initiated by Quark Pharmaceuticals Inc (USA) (<http://quarkpharma.com/?p=12241>).

COMPETING INTERESTS

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank Quark Pharmaceuticals Inc for providing their proprietary siRNA, QPI-1007, targeting caspase-2 (siCASP2) and the control scrambled siRNA (siCNL). This work was funded by a Wellcome Trust project grant no. 092539/Z/10/Z to ZA.

V Vigneswara and Z Ahmed

Neurotrauma Research Group, Institute of Inflammation and Ageing,
College of Medical and Dental Sciences, University of Birmingham,
Birmingham, UK

Correspondence: Z Ahmed (z.ahmed.1@bham.ac.uk)

REFERENCES

- 1 Troy CM et al. *J Neurosci* 2000; **20**: 1386–1392.
- 2 Kisilwa L et al. *Exp Eye Res* 2010; **91**: 739–747.
- 3 Ahmed Z et al. *Cell Death Dis* 2011; **2**: e173.
- 4 Chierzi S et al. *Vision Res* 1998; **38**: 1537–1543.
- 5 Monnier PP et al. *J Neurosci* 2011; **31**: 10494–10505.
- 6 Vigneswara V et al. *PLoS One* 2012; **7**: e53473.
- 7 Vigneswara V et al. *Brain* 2014; **137**: 1656–1675.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016

Supplementary Information accompanies the paper on the Cell Death and Discovery website (<http://www.nature.com/cddiscovery>)